


UNCLASSIFIED

An Affirmative Action/Equal Opportunity Employer

Edited by Jill Warren
Photocomposition by Barbara J. Velarde

NOTICE: Reproduction of this document requires the written consent of the originator, his successor, or higher authority.



DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily reflect those of the United States Government or any agency thereof.

UNCLASSIFIED

~~SECRET~~
UNCLASSIFIED

LOW-YIELD NUCLEAR EXPLOSION CALCULATIONS:
THE 9/22/79 VELA SIGNAL (U)

by

E. M. Jones, R. W. Whitaker,
H. G. Horak, and J. W. Kodis

ABSTRACT (SRD)

DOE
b(1)

Normally, independent confirming evidence is available from other VELA satellites and other sources. Unfortunately, such confirming evidence has not been uncovered for the 9/22/79 event.

DOE
b(1)

In this report we summarize the Vela data, discuss classical interpretations, and present a particular model which, we believe, satisfactorily reproduces the Vela signal.

DOE
b(1)

Thus, our model is consistent with the apparent absence of nuclear debris, the collection of which is required by some analysts for absolute confirmation of an atmospheric detonation.

I. THE VELA DATA

The VELA satellite and its detectors are described elsewhere (e.g. Horak 1980). For our purposes, the salient features of the detection system are:

(
C
C

? DOE
b(1)
b6
b7c

DOE
b1

~~SECRET~~
UNCLASSIFIED

UNCLASSIFIED

**DOE
b(1)**

(min), and the second maximum (2-max) from each of the detectors. The uncertainties are estimated.

- (5) that the instrument response is discrete both in time and in level. The bottoms of adjacent levels are separated by a logarithmic interval.

**DOE
b(1)**

where D is irradiance. The time spacing is somewhat more complex but is basically logarithmic;

(6)

**DOE
b(1)**

The VELA data for the 9/22/79 event are given as irradiance in watts per centimeter squared versus time in milliseconds on the left-hand scale and LD (Level Discriminator) levels on the right-hand scale of Fig. 1. Uncertainties are of the order of one LD level.

Table I lists our best estimates of the times (*t*) and irradiance levels (*I*) at first maximum (1-max), minimum

DUE 5(1)
II. SCALING LAWS FOR ATMOSPHERIC NUCLEAR EXPLOSIONS

Our purpose is to determine the parameters of an atmospheric nuclear explosion that best fit the data

**DOE
b(1)**

UNCLASSIFIED

for each

~~SECRET~~ UNCLASSIFIED

DOE
b(1)

observed by the VELA satellite. As a first step, we use classic scaling laws to estimate the significant parameters of yield and burst height density; then we discuss computer calculations chosen to provide more precise estimates of the explosion parameters.

Numerous sets of scaling laws exist. Here, we have synthesized four adequate scaling laws from discussions by Zing et al. (1974) and Sappenfield (1979).

As we shall see, the time of first maximum and the power ratios are inconsistent with the other observations. Solving the puzzle will concern most of the rest of our study.

DOE
b(1)

DOE
b(1)

The values of the observable quantities ($t_{1,max}$, $t_{2,max}$, and $P_{1,max}P_{2,max}$) have large uncertainties.

Clearly, conventional scaling laws cannot provide a consistent model for the VELA data.

DOE
b(1)

DOE
b(1)

Both Y and ρ are model independent because they are the basic parameters of any calculation. The other three factors can vary dramatically among different models. Let us now consider these other factors and discuss how they can affect the signal.

~~SECRET~~

~~SECRET~~ UNCLASSIFIED

UNCLASSIFIED

significant modification of the signal by the atmosphere or by clouds.

Nuclear weapons with significant neutron output can create an obscuring curtain of "smog" (principally NO_2 , HNO_3 , and O_3). As discussed by Zinn et al. (1974), smog may produce not only a delay in first maximum but also a severe depression of the first pulse by absorption. The minimum and second maximum usually are unaffected by smog because the fireball radius at these times exceeds the distances at which significant neutron deposition occurs.

Perhaps the most dramatic mass effect has been known since the earliest days of atmospheric testing: the apparent yield of a burst at the earth's surface (land or sea) is twice the actual yield. The reason is simply that the surface acts as a nearly perfect reflecting plane. The entire explosion energy acts only in the hemisphere above the surface as if the surface were absent and we were witnessing the spherical expansion of a higher yield explosion. The result is that the radiative/hydrodynamic behavior is identical to that of a free-air burst at twice the actual yield. Among others, Sappenfield (1979) has reviewed the empirical evidence supporting this result.

Cloud cover, atmospheric absorption, and scattering by particulates between the explosion and the satellite can alter the interpretation of the signal. Atmospheric transmission must be considered when deriving radiance at the source from the observed irradiance. However, simple absorption by the atmosphere will not produce dramatic shifts in the timings of events in the signal nor, to first order, will it depress one part of the signal relative to another. However, cloud cover can introduce time smearing caused by photon scattering. Multiple photon scatterings can delay first maximum but should also depress the signal in both peaks. The observed irradiance levels are high enough that the signal could not have suffered much absorption nor scattering beyond that expected for relatively clear maritime air. We do not believe that the data supports the contention that there was any

The hydrodynamic behavior and the power-time curve of this case are illustrated in Figs. 3 and 4, respectively.

UNCLASSIFIED

UNCLASSIFIED
put cap

~~SECRET~~ UNCLASSIFIED

Dec

~~SECRET~~ UNCLASSIFIED

UNCLASSIFIED

Doe
b(3)

" During radial expansion, the fireball brightness depends primarily on the surface area, and the power output increases monotonically. Several bomb masses of air are engulfed during this phase. However, after formation of the air shock at the fireball edge, the brightness depends on the shock speed as well as on the surface area.

Doe
b(3)

Fireball growth is slowed and, more important, areal emission decreases rapidly. The fireball brightness begins to drop when first maximum has been passed.

Doe
b(3)

b(3)

UNCLASSIFIED

~~SECRET~~ UNCLASSIFIED

DOE
b(3)

The rest of this report discusses the effects of mass on the complete optical signal.

III. RADFLO CALCULATIONS

For our present purposes, we have chosen to do calculations in which the nuclear explosive is modeled as a sphere of high-density air at uniform density. The "air bomb" comprises the 10 innermost cells in the calculational grid.

What matters here is the kinetic energy content of the debris, which is largely insensitive to the material composition.

The principal results are tabulated in Tables II and III.

The effect of M and ρ on the times of minimum and of the two maximums are shown in Figs. 7-9. The choice of a time value from a calculation or from data is subjective.

In each calculation, a certain fraction of the explosion energy is deposited as internal energy at $t = 0$ in the central cell.

DOE
b(3)

DOE
b(3)

DOE
b(3)

UNCLASSIFIED

UNCLASSIFIED

X

Doe
b(3)

Doe
b(3)

UNCLASSIFIED

DOE b(3)

UNCLASSIFIED

DOE
b(3)

(3)

The cause of this minimum or its reality is unknown.

DOE
b(3)

UNCLASSIFIED

11

b6 b7
DOE b6 b7
DOE b6 b7

maximum

imums $P_{2-\max}/P_{1-\max}$ appears to have only slight sensitivity to mass and ambient density.

Zian et al. (1974) show a strong $t_{1-\max}$ dependence on density. Their result is confirmed by our calculations.

The ratio $P_{2-\max}/P_{1-\max}$ is a strong function of mass, yield, and density.

b6 b7 (11)

These scaling laws [Eqs. (6)-(8)] can be used to refine our estimates of the explosion parameters. The very weak density dependence and the large timing uncertainty that we cannot derive a dependable density

b6 b7
DOE b6 b7

We return to a discussion of burst height effects in Sec. IV.

We can now derive a yield and mass from our scaling laws.

IV. A MODEL FOR THE 9/22/79 BURST (ALERT 747)

b6 b7

b6 b7
DOE b6 b7


Figure 9 shows the band of permitted values in the (Y,M) plane. The $t_{1-\max}$ scaling law [Eq. (6)] seems to give a slight overestimate of $t_{1-\max}$ at high masses.

As a final check on the explosion parameters, we note their effect on the power ratios $P_{2-\max}/P_{1-\max}$ and $P_{2-\max}/P_{\min}$ as shown in Figs. 10-12. The ratio of max-

b6 b7
DOE b6 b7

The predicted power-time curve is shown in Fig. 12. This is a time-averaged curve.

UNCLASSIFIED

 UNCLASSIFIED

DOE
b(3)

J
13

 UNCLASSIFIED

~~UNCLASSIFIED~~

b(3)

$$\bar{P} = \frac{1}{dt} \int_{cycle} P dt \quad (13)$$

V. PERTINENT DATA FROM ATMOSPHERIC NUCLEAR TESTS

We have constructed what we believe to be a plausible model for a low-yield nuclear explosion that could have

b(3) (15)

The agreement is satisfactory.

After the shock emerges from the bomb and begins to sweep up ambient air, the power-time curve shows dramatic oscillations produced by the entrainment of individual calculational cells. When the shock first encounters an ambient cell, the air temperature in the cell rises, causing the opacity to rise and thereby decreasing fireball brightness. Eventually, the newly entrained cell is heated enough that it starts radiating, and the fireball brightness increases until the cycle is repeated with the entrainment of the next cell outward. Interpretation of the calculated power levels during the first pulse is uncertain. We have chosen to show the power levels averaged over each cycle as

The calculated amplitude of the bump at high mass is well below the data uncertainties.

At later times, physical mechanisms responsible for producing the power-time curve variations are well understood. The interested reader may consult Zinn (1973), Zinn et al. (1974), Brode (1968), or Glasstone (1964) for discussions of phenomena at t_{max} and beyond.

We have plotted our best interpretation of the VELA data to produce the power outputs in Fig. 12.

(12)

b(3)

b(3)

b(3)



UNCLASSIFIED

infuse the observed Alert 747 signal.

b6 b7c
In this section, we examine the
available data for t_{rise} from atmospheric nuclear tests.

The available t_{rise} data are plotted as a function of
yield in Fig. 12. We note that for many events of the US
atmospheric test program, data pertinent to the first
maximum are not available.

b6 b7c
Doc 6(3) 4/12

b6 b7c
6(3)

However, the fit after about ms
is quite satisfactory

b6 b7c
6(3)

*Personal communication to authors.

b6 b7c
6(3)

UNCLASSIFIED¹⁵

~~UNCLASSIFIED~~

b6
b7c
b8c

GEORGE was detonated atop a tower 61 m above Eniwetok on May 9, 1951.

b6
b7c

Two features in the explosion phenomenology are of interest. First, the fireball expansion was very asymmetric during the early phases; second, independent determination of the yield by hydrodynamic and radiochemical techniques were widely discrepant.

b6
b7c

The other part of the fireball may have grown radiatively. Clearly, calculational studies in two or three dimensions will be required to provide a convincing explanation.

At later times, the expansion becomes more symmetric and, at minimum, this fireball has prominent spots.

b6
b7c

16



b6
b7c

b6
b7c

The hydrodynamic yield is determined by comparing the radius-time data with the expectations of classic blast wave theory. In particular, during the period immediately preceding minimum, the quality

b6
b7c

(16)

is virtually constant. In Fig. 16, we show the ϕ histories obtained from four RADFLO calculations compared



~~UNCLASSIFIED~~

DoE b(3)

UNCLASSIFIED

DoE
b(3)

VI. CAVEATS

DoE
b(1)

with the GEORGE data. These data were derived by visually estimating a "best-circle" fit to the fireball shape.

Nonetheless, we recognize that we have made certain assumptions that, if proved wrong, may alter our conclusions.

DoE
b(1)

- (2) We have used dense air to model the weapon vapor.

DoE
b(3)

DoE b(3)

UNCLASSIFIED¹⁷

UNCLASSIFIED

US DOE, NERI
WASH.

(3)

DOE
b(3)

We have been unable to secure experimental confirmation of a $t_{1-\alpha}$ delay caused by mass.

(4)

DOE
b(1)
b(3)

The possibility remains that other models can produce good fits to the data. Proponents of such models must demonstrate plausibility.

VII. SUMMARY

ACKNOWLEDGMENTS

We gratefully acknowledge valuable discussion with H. Hoerlin, J. Zinn, and J. Malik of Los Alamos National Laboratory and G. Mauch of Sandia National Laboratories.

REFERENCES

- Brode, H. L., "Numerical Solutions of Spherical Blast Waves," *J. App. Phys.* 26, 766 (1955).
- Brode, H. L., "Review of Nuclear Weapons Effects," *Ann. Rev. Nuc. Sci.* 18, 153 (1968).
- Colvin, J. D. and N. P. Gow, "Fireball Spots: A Comprehensive Study of the Photographic Data (U)," EG&G Technical Report EGG 1183-5086 (SFRD) (August 1978).
- Glasstone, S. (Ed), *The Effects of Nuclear Weapons - Revised Edition (U)* (Washington: U.S. Atomic Energy Commission 1964).
- Horak, H. G., "Vela Event Alert 747 (U)," Los Alamos Scientific Laboratory Report LA-8364-MS (SRD) (May 1980).
- Horak, H. G. and J. W. Kodis, "RADFLO: A User's Manual (U)," Los Alamos National Laboratory Report LA-9245-M (U) submitted for publication.
- Sappenfield, D. S., "The Surface Burst Correction Factor for Barograph Scaling (U)," Air Force Technical Applications Center Topical Report AFTAC-TR-79-12 (SRD) (March 1979).
- Sedov, L. I., *Similarity and Dimensional Methods in Mechanics* (New York: Academic Press, 1959).
- Taylor, G. I., "The Formation of a Blast Wave by a Very Intense Explosion. I. Theoretical Discussion," *Proc. Roy. Soc. A201*, 159 (1950).
- Zinn, J., "A Finite Difference Scheme for Time-Dependent Spherical Radiation Hydrodynamics Problems," *J. Comp. Phys.* 13, 569 (1973).
- Zinn, J., J. W. Kodis, and C. D. Sutherland, "Status Report on Fireball Radiation Hydrodynamics Computations (U)," Los Alamos Scientific Laboratory Report LA-5591-SR (SFRD) (May 1974).

*Private communication to authors.

UNCLASSIFIED

UNCLASSIFIED

US DOE, Headquarters Library, Reports Section,
Washington, DC
Manager, ALO, Albuquerque, New Mexico
Lawrence Livermore National Laboratory,
Livermore, California
Sandia National Laboratories, Albuquerque, New Mexico
Military Liaison Committee, Washington, DC
Headquarters, Defense Nuclear Agency, Washington, DC
Defense Nuclear Agency Field Command,
Kirtland AFB, New Mexico

Standard Distribution

Copy No.		Copy No.
1-3	Commander, Naval Sea Systems Command, Washington, DC	13
4	Director, Air Force Weapons Laboratory, Kirtland AFB, New Mexico	14-16
5-6	US DOE, Technical Information Center, Oak Ridge, Tennessee	17
7	Attn: Weapon Data Index	
8	Los Alamos National Laboratory Report Library	
9-10		
11-12		

Special Distribution

Central Intelligence Agency
Washington, DC
Attn: Chief, OSWR/NED
For: Frank P. Burrowski, JAEIC
Donald M. Braisted, CNED (2 exs.)
LT. GEN. Harold C. Donnelly, JAEIC
LTC. James Farga, DAMI-FIT
Kenneth Gilbert, NSA/AOS
LT. COL. Houston Hawkins,
DIA(DT-IE)
Dr. John S. Ingley, OSWR
Dr. Gerald W. Johnson
Max A. Koontz, NSC
Robert Morrison, ACDA
Dr. S. A. Newton, Jr., JAEIC
Louis H. Roddis, Jr., JAEIC
Robert Rubinstein, DOE
Jeffry Siegel, JAEIC
Dr. Gerald F. Tape, JAEIC
CAPT. Raymond Vikus, AF/INET
VADM Steven A. White

Lawrence Livermore National Laboratory
Livermore, California
Attn: C. Alonso
R. Klein
Dr. Roger E. Batzel
Dr. Edward Teller

Sandia National Laboratories
Albuquerque, New Mexico
Attn: Mail Services Section
For: Glena Fowler
A. Chabai
G. Meuth
D. Thornborough

EG&G, Inc.
Los Alamos, New Mexico
Attn: Document Control
For: D. Wright
J. Colvin
L. Rumber

Copy No.		Copy No.
65-82	Defense Nuclear Agency, Headquarters Department of Defense Washington, DC Director Attn: PAAD-3 For: C. Fitz L. Wittwer G. Sevin C. Knowles	94-97
83-86	Air Force Technical Applications Center Patrick Air Force Base, Florida AFTAC/TD Attn: Dr. Gerard M. Laine	98
87-90	Air Force Weapons Laboratory Kirtland Air Force Base, New Mexico Attn: SUL, Mrs. Georgiana Hilfner For: MAJ. Gasong	
91-93	Union Carbide Corporation Y-12 Plant Nuclear Division Oak Ridge, Tennessee Attn: Y-12 Plant Records Department For: Paul R. Vanstrum	101
	Harold M. Agnew, Consultant c/o General Atomic Company La Jolla, California	102
	Walter G. Chestnut, Consultant Menlo Park, California	103
	W. E. Ogle, Consultant Energy Systems, Inc. Anchorage, Alaska Attn: W. E. Ogle	104

UNCLASSIFIED

UNCLASSIFIED

now now

now now UNCLASSIFIED